

This article was downloaded by: [Nevsehir Universitesi]

On: 11 June 2015, At: 03:36

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Journal of Biological Dynamics

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tjbd20>

### Global behaviour of a predator-prey like model with piecewise constant arguments

Senol Kartal<sup>a</sup> & Fuat Gurcan<sup>bc</sup>

<sup>a</sup> Department of Mathematics, Faculty of Education, Nevsehir Haci Bektas Veli University, Nevsehir 50300, Turkey

<sup>b</sup> Faculty of Engineering and Natural Sciences, International University of Sarajevo, Hrasnicka cesta 15, 71000 Sarajevo, BIH

<sup>c</sup> Department of Mathematics, Faculty of Science, Erciyes University, Kayseri 38039, Turkey

Published online: 04 Jun 2015.



CrossMark

[Click for updates](#)

To cite this article: Senol Kartal & Fuat Gurcan (2015) Global behaviour of a predator-prey like model with piecewise constant arguments, *Journal of Biological Dynamics*, 9:1, 159-171, DOI: [10.1080/17513758.2015.1049225](https://doi.org/10.1080/17513758.2015.1049225)

To link to this article: <http://dx.doi.org/10.1080/17513758.2015.1049225>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Versions of published Taylor & Francis and Routledge Open articles and Taylor & Francis and Routledge Open Select articles posted to institutional or subject repositories or any other third-party website are without warranty from Taylor & Francis of any kind, either expressed or implied, including, but not limited to, warranties of merchantability, fitness for a particular purpose, or non-infringement. Any opinions and views expressed in this article are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor & Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

It is essential that you check the license status of any given Open and Open Select article to confirm conditions of access and use.

# Global behaviour of a predator–prey like model with piecewise constant arguments

Senol Kartal<sup>a\*</sup> and Fuat Gurcan<sup>b,c</sup>

<sup>a</sup>Department of Mathematics, Faculty of Education, Nevsehir Haci Bektas Veli University, Nevsehir 50300, Turkey; <sup>b</sup>Faculty of Engineering and Natural Sciences, International University of Sarajevo, Hrasnicka cesta 15, 71000 Sarajevo, BIH; <sup>c</sup>Department of Mathematics, Faculty of Science, Erciyes University, Kayseri 38039, Turkey

(Received 10 November 2014; accepted 4 May 2015)

The present study deals with the analysis of a predator–prey like model consisting of system of differential equations with piecewise constant arguments. A solution of the system with piecewise constant arguments leads to a system of difference equations which is examined to study boundedness, local and global asymptotic behaviour of the positive solutions. Using Schur–Cohn criterion and a Lyapunov function, we derive sufficient conditions under which the positive equilibrium point is local and global asymptotically stable. Moreover, we show numerically that periodic solutions arise as a consequence of Neimark–Sacker bifurcation of a limit cycle.

**Keywords:** piecewise constant arguments; difference equation; stability; bifurcation

*AMS Subject Classification:* 39A10; 39A28; 39A30

## 1. Introduction

Recently, there has been great interest in studying differential equations with piecewise constant arguments because of the wide application of these equations in biology, engineering and other fields. Many authors have analysed various types of population models based on logistic equations with piecewise constant arguments and have obtained theoretical results on oscillations or chaotic behaviour [2,4–6,8–14,15,16]. The simplest model was proposed by May [9] and May and Oster [10] who obtained that the model have chaotic behaviour for certain parameters. On the other hand, several authors [8,11,12,15,16] have investigated a more general logistic equation with piecewise constant arguments

$$\frac{dx(t)}{dt} = rx(t)(1 - ax(t) - b \sum_{j=0}^m c_j x([t-j])), \quad t \geq 0. \quad (1)$$

\*Corresponding author. Email: [senol.kartal@nevsehir.edu.tr](mailto:senol.kartal@nevsehir.edu.tr)  
Author Email: [gurcan@erciyes.edu.tr](mailto:gurcan@erciyes.edu.tr)

Liu and Gopalsamy [8] have showed that for certain special cases, solutions of equation (1) can have chaotic behaviour through period doubling bifurcations. Muroya [12] has improved contractivity conditions for the positive equilibrium point of this equation.

Following these works, Gurcan and Bozkurt [5] have studied global stability and boundedness character of the positive solutions of the differential equation

$$\frac{dx(t)}{dt} = rx(t)(1 - \alpha x(t) - \beta_0 x([t]) - \beta_1 x([t-1])). \quad (2)$$

By using this equation, Ozturk *et al.* [14] have modelled a population density of a bacteria species in a microcosm. A more general case of equation (2) has been considered by Ozturk and Bozkurt [13] as the following;

$$\frac{dx(t)}{dt} = x(t)(r(1 - \alpha x(t) - \beta_0 x([t]) - \beta_1 x([t-1])) + \gamma_1 x([t]) + \gamma_2 x([t-1])). \quad (3)$$

They have investigated stability and oscillatory characteristics of difference solutions of the equation. Equation (3) has also been used for modelling an early brain tumour growth in [2]. The stability analysis of the model shows that increase in the tumour growth rate decreases the local stability of the positive equilibrium point. Another mathematical model for tumour growth under the immune activity has been constructed Banerjee and Sarkar [1] such as

$$\begin{aligned} \frac{dM}{dt} &= r_1 M \left(1 - \frac{M}{k_1}\right) - \alpha_1 MN, \\ \frac{dN}{dt} &= \beta NZ(t - \tau) - d_1 N - \alpha_2 MN, \\ \frac{dZ}{dt} &= r_2 Z \left(1 - \frac{Z}{k_2}\right) - \beta NZ(t - \tau), \end{aligned} \quad (4)$$

where  $M(t)$ ,  $N(t)$ ,  $Z(t)$  represent the number of tumour, hunting and resting cells, respectively. The model consists of delay differential equations which often arise in biological systems. Since analysis of these equations is more complicated than ordinary differential equations, numerical approach may be needed for delay differential equations. In study [3], Cooke and Györi have showed that differential equations with piecewise constant arguments can be used to obtain approximate solution to delay differential equations that contain discrete delays.

In the present paper, we have modified model (4) by adding piecewise constant arguments such as

$$\begin{aligned} \frac{dM}{dt} &= r_1 M(t) \left(1 - \frac{M(t)}{k_1}\right) - \alpha_1 M(t)N([t]), \\ \frac{dN}{dt} &= \beta N(t)Z([t]) - d_1 N(t) - \alpha_2 M([t])N(t), \\ \frac{dZ}{dt} &= r_2 Z(t) \left(1 - \frac{Z(t)}{k_2}\right) - \beta N([t])Z(t), \end{aligned} \quad (5)$$

where  $[t]$  denotes the integer part of  $t \in [0, \infty)$  and all these parameters are positive. The model includes both discrete and continuous time for tumour, hunting and resting cells because tumour cells have different dynamics which can be described by using both differential and difference equations. Here,  $M(t)$ ,  $N(t)$  and  $Z(t)$  represent population density of tumour, hunting and resting cells, respectively. The parameters  $r_1$  and  $k_1$  represent the growth rate and the maximum carrying capacity of tumour cells, respectively.  $r_2$  is the growth rate,  $k_2$  is the maximum carrying capacity

of resting cells and  $d_1$  is the natural death of hunting cell. The parameter  $\alpha_1$  denotes decay rate of tumour cells by hunting cells,  $\alpha_2$  is decay rate of hunting cells by tumour cells and  $\beta$  is conversion rate from resting to hunting cells. Most of the parameter values are taken from the [1] in terms of consistency with the biological facts. In Section 2, we investigate boundedness, local and global behaviour of the positive solutions of the system by using the method of reduction to discrete equations. In Section 3, we study periodic solution of the system through Neimark-Sacker bifurcation.

## 2. Boundedness, local and global stability analysis

In this section, by integrating of system (5) we first obtain a solution and later discuss the boundedness and the local asymptotic stability of system (7).

We can rewrite system (5) on an interval of the form  $t \in [n, n + 1)$  as follows:

$$\begin{aligned}\frac{dM}{dt} &= r_1 M(t) \left(1 - \frac{M(t)}{k_1}\right) - \alpha_1 M(t) N(n), \\ \frac{dN}{dt} &= \beta N(t) Z(n) - d_1 N(t) - \alpha_2 M(n) N(t), \\ \frac{dZ}{dt} &= r_2 Z(t) \left(1 - \frac{Z(t)}{k_2}\right) - \beta N(n) Z(t).\end{aligned}\tag{6}$$

By solving each equations in the system (6) with respect to  $t$  on  $[n, t)$  and letting  $t \rightarrow n + 1$ , we obtain a system of difference equations

$$\begin{aligned}M(n+1) &= \frac{M(n)(r_1 - \alpha_1 N(n))}{r_1 K_1 M(n) + (r_1 - \alpha_1 N(n) - r_1 K_1 M(n)) \exp(-(r_1 - \alpha_1 N(n)))}, \\ N(n+1) &= N(n) \exp(\beta Z(n) - d_1 - \alpha_2 M(n)), \\ Z(n+1) &= \frac{Z(n)(r_2 - \beta N(n))}{r_2 K_2 Z(n) + (r_2 - \beta N(n) - r_2 K_2 Z(n)) \exp(-(r_2 - \beta N(n)))},\end{aligned}\tag{7}$$

where  $1/k_1 = K_1$ ,  $1/k_2 = K_2$ . Thus, we obtain discrete analogue of system (5) as a system of difference equation which reveals the rich dynamical characteristics and the asymptotic behaviour of the dynamical system (5). Now, we can discuss the boundedness of solutions of the system in the following theorem.

**THEOREM 2.1** *Let  $\{M(n), N(n), Z(n)\}_{n=-1}^{\infty}$  be a positive solution of system (7); then*

$$0 \leq M(n) \leq \frac{\exp(r_1)}{K_1(\exp(r_1) - 1)} \quad \text{and} \quad 0 \leq Z(n) \leq \frac{\exp(r_2)}{K_2(\exp(r_2) - 1)}.$$

*In addition, if  $\beta Z(n) - d_1 - \alpha_2 M(n) < 0$ , then  $0 \leq N(n) \leq N(0)$ .*

*Proof* It is easy to see that system (5) can be written on an interval of the form  $t \in [n, n + 1)$  as follows:

$$\begin{aligned} M(t) &= M(n) \exp\left(\int_n^t (r_1(1 - M(s)K_1) - \alpha_1 N(n)) ds\right), \\ N(t) &= N(n) \exp((\beta Z(n) - d_1 - \alpha_2 M(n))(t - n)), \\ Z(t) &= Z(n) \exp\left(\int_n^t (r_2(1 - Z(s)K_2) - \beta N(n)) ds\right). \end{aligned}$$

It is clear that if  $M(0) > 0, N(0) > 0$  and  $Z(0) > 0$ , then  $M(t) > 0, N(t) > 0$  and  $Z(t) > 0$  for  $t > 0$ . This implies that we have positive solutions of Equation (5) for positive initial conditions.

From the first equation in system (7), we have

$$\begin{aligned} M(n+1) &= \frac{M(n)(r_1 - \alpha_1 N(n)) \exp(r_1 - \alpha_1 N(n))}{r_1 - \alpha_1 N(n) + r_1 K_1 M(n) (\exp(r_1 - \alpha_1 N(n)) - 1)} \\ &\leq \frac{M(n)(r_1 - \alpha_1 N(n)) \exp(r_1 - \alpha_1 N(n))}{r_1 K_1 M(n) (\exp(r_1 - \alpha_1 N(n)) - 1)} \\ &= \frac{(r_1 - \alpha_1 N(n)) \exp(r_1 - \alpha_1 N(n))}{r_1 K_1 (\exp(r_1 - \alpha_1 N(n)) - 1)} \\ &\leq \frac{r_1 \exp(r_1)}{r_1 K_1 (\exp(r_1) - 1)} \\ &= \frac{\exp(r_1)}{K_1 (\exp(r_1) - 1)}. \end{aligned}$$

Additionally, It can be easily shown that  $Z(n) \leq \exp(r_2)/K_2(\exp(r_2) - 1)$ . Now, we consider the second equation in system (7). Under the condition  $\beta Z(n) - d_1 - \alpha_2 M(n) < 0$ , we get

$$N(n+1) = N(n) \exp(\beta Z(n) - d_1 - \alpha_2 M(n)) \leq N(n).$$

This completes the proof. ■

To analyse global behaviour of the difference system, we need to determine the positive equilibrium point. Let

$$\begin{aligned} f_1(M(n), N(n), Z(n)) &= \frac{M(n)(r_1 - \alpha_1 N(n))}{r_1 K_1 M(n) + (r_1 - \alpha_1 N(n) - r_1 K_1 M(n)) \exp(-(r_1 - \alpha_1 N(n)))}, \\ f_2(M(n), N(n), Z(n)) &= N(n) \exp(\beta Z(n) - d_1 - \alpha_2 M(n)), \\ f_3(M(n), N(n), Z(n)) &= \frac{Z(n)(r_2 - \beta N(n))}{r_2 K_2 Z(n) + (r_2 - \beta N(n) - r_2 K_2 Z(n)) \exp(-(r_2 - \beta N(n)))}. \end{aligned}$$

Thus, the positive equilibrium point of system (7) or fixed point of the vector map  $\mathbf{F} = (f_1, f_2, f_3)$  can be obtained from the solution of the system

$$\begin{aligned} \bar{M} &= f_1(\bar{M}, \bar{N}, \bar{Z}), \\ \bar{N} &= f_2(\bar{M}, \bar{N}, \bar{Z}), \\ \bar{Z} &= f_3(\bar{M}, \bar{N}, \bar{Z}), \end{aligned}$$

as  $\bar{E} = (\bar{M}, \bar{N}, \bar{Z})$  where

$$\bar{M} = \frac{\beta^2 r_1 - \beta r_2 \alpha_1 + d_1 K_2 r_2 \alpha_1}{\beta^2 K_1 r_1 - K_2 r_2 \alpha_1 \alpha_2}, \quad \bar{N} = \frac{r_1 r_2 (\beta K_1 - d_1 K_1 K_2 - K_2 \alpha_2)}{\beta^2 K_1 r_1 - K_2 r_2 \alpha_1 \alpha_2},$$

and

$$\bar{Z} = \frac{\beta d_1 K_1 r_1 + \beta r_1 \alpha_2 - r_2 \alpha_1 \alpha_2}{\beta^2 K_1 r_1 - K_2 r_2 \alpha_1 \alpha_2}.$$

For  $\bar{M} > 0, \bar{N} > 0$  and  $\bar{Z} > 0$ , we must hold conditions

$$\beta^2 r_1 - \beta r_2 \alpha_1 + d_1 K_2 r_2 \alpha_1 > 0, \tag{8}$$

$$\beta^2 K_1 r_1 - K_2 r_2 \alpha_1 \alpha_2 > 0, \tag{9}$$

$$\beta K_1 - d_1 K_1 K_2 - K_2 \alpha_2 > 0, \tag{10}$$

$$\beta d_1 K_1 r_1 + \beta r_1 \alpha_2 - r_2 \alpha_1 \alpha_2 > 0. \tag{11}$$

By analysing conditions (8)–(11) with respect to  $\alpha_1$  and  $\beta$ , we can obtain the inequalities

$$\alpha_1 < \frac{\beta^2 r_1}{\beta r_2 - d_1 K_2 r_2} \quad \text{and} \quad \beta > \frac{d_1 K_1 K_2 + K_2 \alpha_2}{K_1}. \tag{12}$$

The Jacobian matrix of map  $\mathbf{F} = (f_1, f_2, f_3)$  at positive equilibrium point  $\bar{E}$  is the matrix

$$\mathbf{J}_F(\bar{E}) = \begin{pmatrix} \exp(-K_1 r_1 \bar{M}) & -\frac{(1 - \exp(-K_1 r_1 \bar{M}))\alpha_1}{K_1 r_1} & 0 \\ -\alpha_2 \bar{N} & 1 & \beta \bar{N} \\ 0 & -\frac{(1 - \exp(-K_2 r_2 \bar{Z}))\beta}{K_2 r_2} & \exp(-K_2 r_2 \bar{Z}) \end{pmatrix}.$$

Under the assumption

$$\exp(-K_1 r_1 \bar{M}) = \exp(-K_2 r_2 \bar{Z}), \tag{13}$$

the characteristic polynomial of the matrix  $\mathbf{J}_F(\bar{E})$  at the positive equilibrium point is

$$p_1(\lambda) = (\exp(-K_1 r_1 \bar{M}) - \lambda) \left( (1 - \lambda)(\exp(-K_1 r_1 \bar{M}) - \lambda) + \frac{\bar{N}(1 - \exp(-K_1 r_1 \bar{M}))(\beta^2 K_1 r_1 - K_2 r_2 \alpha_1 \alpha_2)}{K_1 r_1 K_2 r_2} \right).$$

From the first factor in this equation, an eigenvalue is computed as  $\lambda_1 = \exp(-K_1 r_1 \bar{M}) < 1$ . Solving Equation (13), we have

$$d_1 = \frac{(\beta r_1 - r_2 \alpha_1)(\beta K_1 r_1 - K_2 r_2 \alpha_2)}{K_1 r_1 K_2 r_2 (\beta - \alpha_1)}. \tag{14}$$

Thus, the characteristic polynomial is reduced to a second-order equation

$$p_2(\lambda) = \lambda^2 + \lambda(-1 - \exp(-K_1 r_1 \bar{M})) + \exp(-K_1 r_1 \bar{M}) + \frac{\bar{N}(1 - \exp(-K_1 r_1 \bar{M}))(\beta^2 K_1 r_1 - K_2 r_2 \alpha_1 \alpha_2)}{K_1 r_1 K_2 r_2}.$$

To determine stability conditions of discrete system (7) through the characteristic equation  $p_2(\lambda)$ , we can use Schur–Chon criterion.

**THEOREM 2.2** Let  $\bar{E} = (\bar{M}, \bar{N}, \bar{Z})$  be the positive equilibrium point of system (7) and

$$d_1 = \frac{(\beta r_1 - r_2 \alpha_1)(\beta K_1 r_1 - K_2 r_2 \alpha_2)}{K_1 r_1 K_2 r_2 (\beta - \alpha_1)}.$$

If

$$\alpha_1 < \frac{\beta^2 r_1}{\beta r_2 - d_1 K_2 r_2} \quad \text{and} \quad \frac{d_1 K_1 K_2 + K_2 \alpha_2}{K_1} < \beta < \frac{K_1 K_2 + d_1 K_1 K_2 + K_2 \alpha_2}{K_1},$$

then  $\bar{E}$  is local asymptotically stable.

*Proof* By the Schur–Cohn criterion, we get that  $\bar{E}$  is local asymptotically stable if and only if

$$| -1 - \exp(-K_1 r_1 \bar{M}) | < 1 + \exp(-K_1 r_1 \bar{M}) + \frac{\bar{N}(1 - \exp(-K_1 r_1 \bar{M}))(\beta^2 K_1 r_1 - K_2 r_2 \alpha_1 \alpha_2)}{K_1 r_1 K_2 r_2} \quad (15)$$

and

$$1 + \exp(-K_1 r_1 \bar{M}) + \frac{\bar{N}(1 - \exp(-K_1 r_1 \bar{M}))(\beta^2 K_1 r_1 - K_2 r_2 \alpha_1 \alpha_2)}{K_1 r_1 K_2 r_2} < 2. \quad (16)$$

It is easy to see that Equation (15) always exists. From Equation (16), we have

$$\frac{\bar{N}(1 - \exp(-K_1 r_1 \bar{M}))(\beta^2 K_1 r_1 - K_2 r_2 \alpha_1 \alpha_2) + K_1 r_1 K_2 r_2 \exp(-K_1 r_1 \bar{M})}{K_1 r_1 K_2 r_2} < 1$$

which reveal that

$$\beta < \frac{K_1 K_2 + d_1 K_1 K_2 + K_2 \alpha_2}{K_1}.$$

This completes the proof. ■

*Example 2.3* In this example, all parameter values are taken in [1] as  $r_1 = 0.18 \text{ day}^{-1}$ ,  $r_2 = 0.0245 \text{ day}^{-1}$ ,  $k_1 = 5 \times 10^6 \text{ cells}$ ,  $k_2 = 1 \times 10^7 \text{ cells}$ ,  $\beta = 6.2 \times 10^{-9} \text{ cells}^{-1} \text{ day}^{-1}$ ,  $\alpha_2 = 3.422 \times 10^{-10} \text{ cells}^{-1} \text{ day}^{-1}$ ,  $d_1 = 0.0412 \text{ day}^{-1}$ , except  $\alpha_1$ . It can be seen that under the conditions given in Theorem 2.2 and using initial conditions  $M(0) = 4.53941 \times 10^5$ ,  $N(0) = 1.3158 \times 10^6$  and  $Z(0) = 6.67022 \times 10^6$ , the equilibrium point  $\bar{E} = (\bar{M}, \bar{N}, \bar{Z}) = (4.53941 \times 10^5, 1.3158 \times 10^6, 6.67022 \times 10^6)$  is local asymptotically stable where blue, red and black graphs represent  $M(n)$ ,  $N(n)$  and  $Z(n)$  population densities, respectively (see Figure 1).

Using the parameter values given in Example 2.3, positive equilibrium point  $\bar{E} = (4.53941 \times 10^5, 1.3158 \times 10^6, 6.67022 \times 10^6)$  is obtained under the conditions  $\beta > 4.2911 \times 10^{-9}$  and  $\alpha_1 < 1.35777 \times 10^{-7}$  which are exactly the same as in those of Banerjee and Sarkar [1]. In addition, our stability results can be compared numerically with that of work [1]. Although in their delayed system local stability condition on parameter  $\beta$  is  $\beta > 4.2911 \times 10^{-9}$ , we have  $4.2911 \times 10^{-9} < \beta < 1.0429 \times 10^{-7}$  which is obtained from Theorem 2.2. In addition at the present study, the condition on  $\alpha_1$  is  $\alpha_1 < 1.35777 \times 10^{-7}$ , but this condition is determined as  $\alpha_1 < 1.26004 \times 10^{-7}$  in study [1] under the set of our parameter values. These results indicate that there is a little differences between their and our stability conditions.






Figure 1. Graph of the iteration solution of  $M(n), N(n)$  and  $Z(n)$ , where  $\alpha = 1.24379 \times 10^{-7}$ .

**THEOREM 2.4** Let  $A_1 = r_1 - \alpha_1 N(n), A_2 = r_2 - \beta N(n)$  and  $A_3 = \beta Z(n) - d_1 - \alpha_2 M(n)$ . Suppose that the conditions of Theorem 2.2 hold and

$$(a_1) N(n) < \frac{r_1}{\alpha_1} \quad \text{and} \quad \bar{M} < \frac{A_1(1 + \exp(-A_1))}{2r_1 K_1 \exp(-A_1)} \quad \text{for } M(n) \in \left(0, \frac{2\bar{M} \exp(-A_1)}{1 + \exp(-A_1)}\right),$$

$$(a_2) N(n) > \frac{r_1}{\alpha_1} \quad \text{and} \quad \bar{M} > \frac{-A_1(1 + \exp(-A_1))}{2r_1 K_1 (\exp(-A_1) - 1)} \quad \text{for } M(n) \in \left(\frac{2\bar{M} \exp(-A_1)}{1 + \exp(-A_1)}, 2\bar{M}\right),$$

$$(a_3) N(n) > \frac{r_1}{\alpha_1} \quad \text{for } M(n) \in (2\bar{M}, \infty),$$

$$(b_1) A_3 > 0 \quad \text{for } N(n) \in \left(0, \frac{2\bar{N}}{1 + \exp(A_3)}\right),$$

$$(b_2) A_3 < 0 \quad \text{for } N(n) \in \left(\frac{2\bar{N}}{1 + \exp(A_3)}, \infty\right),$$

$$(c_1) N(n) < \frac{r_2}{\beta} \quad \text{and} \quad \bar{Z} < \frac{A_2(1 + \exp(-A_2))}{2r_2 K_2 \exp(-A_2)} \quad \text{for } Z(n) \in \left(0, \frac{2\bar{Z} \exp(-A_2)}{1 + \exp(-A_2)}\right),$$

$$(c_2) N(n) > \frac{r_2}{\beta} \quad \text{and} \quad \bar{Z} > \frac{-A_2(1 + \exp(-A_2))}{2r_2 K_2 (\exp(-A_2) - 1)} \quad \text{for } Z(n) \in \left(\frac{2\bar{Z} \exp(-A_2)}{1 + \exp(-A_2)}, 2\bar{Z}\right),$$

$$(c_3) N(n) > \frac{r_2}{\beta} \quad \text{for } Z(n) \in (2\bar{Z}, \infty).$$

Then the positive equilibrium point of system (7) is global asymptotically stable.

*Proof* Let  $\bar{E} = (\bar{M}, \bar{N}, \bar{Z})$  is positive equilibrium point system (7) and we consider a Lyapunov function  $V(n)$  defined by

$$V(n) = (E(n) - \bar{E})^2, \quad n = 0, 1, 2, \dots$$




Figure 2. Graph of the iteration solution of  $M(n)$ ,  $N(n)$  and  $Z(n)$ , where  $\alpha = 1.24379 \times 10^{-7}$ . Parameter values are taken Example 2.3.

The change along the solutions of the system is

$$\Delta V(n) = V(n+1) - V(n) = (E(n+1) - E(n))(E(n+1) + E(n) - 2\bar{E}).$$

From the first equation in (7), we get

$$\begin{aligned} \Delta V_1(n) &= (M(n+1) - M(n))(M(n+1) + M(n) - 2\bar{M}) \\ &= M(n)((A_1 - r_1 K_1 M(n))(1 - \exp(-A_1))(A_1(M(n) - \bar{M}) \exp(-A_1) \\ &\quad + \exp(-A_1)(M(n) - \bar{M})) + r_1 K_1 M(n)(M(n) - 2\bar{M})(1 - \exp(-A_1))). \end{aligned}$$

For each assumption  $(a_1)$ ,  $(a_2)$  and  $(a_3)$ , we have  $\Delta V_1(n) < 0$  which implies  $\lim_{n \rightarrow \infty} M(n) = \bar{M}$ . Additionally, we hold

$$\begin{aligned} \Delta V_2(n) &= (N(n+1) - N(n))(N(n+1) + N(n) - 2\bar{N}) \\ &= N(n)(\exp(A_3) - 1)(N(n) \exp(A_3) + N(n) - 2\bar{N}) < 0 \end{aligned}$$

for each assumption  $(b_1)$  and  $(b_2)$  which follows  $\lim_{n \rightarrow \infty} N(n) = \bar{N}$ . Similarly, it can be shown that  $\Delta V_3(n) = (Z(n+1) - Z(n))(Z(n+1) + Z(n) - 2\bar{Z}) < 0$  under the assumptions  $(c_1)$ ,  $(c_2)$  and  $(c_3)$ . As a result, we obtain  $\Delta V(n) = (\Delta V_1(n), \Delta V_2(n), \Delta V_3(n)) < 0$ . ■

*Example 2.5* Considering conditions of Theorem 2.4, we can use the parameter values in Example 2.3 and initial conditions  $M(0) = 1.53 \times 10^5$ ,  $N(0) = 2.31 \times 10^5$  and  $Z(0) = 3.67 \times 10^6$ . The graph of the first 3000 iterations of system (7) is given in Figure 2, where blue, red and black graphs represent  $M(n)$ ,  $N(n)$  and  $Z(n)$  population densities, respectively. It can be shown that under the conditions given in Theorem 2.4 the positive equilibrium point is global asymptotically stable.

### 3. Neimark-Sacker bifurcation analysis

The Neimark-Sacker bifurcation is extremely important in the context of discrete biological models, where one observes periodic solutions corresponding to a closed invariant curve (that is a




Figure 3. Graph of the  $(\beta, |\lambda|)$ . Parameter set is taken from Example 2.3.

limit cycle) in the phase space. For this bifurcation, characteristic equation has a pair of complex conjugate eigenvalues on the unit circle and all other eigenvalues are inside the circle.

To study Neimark-Sacker bifurcation as in the work of Banerjee and Sarkar [1], we choose parameter  $\beta$  as a bifurcation parameter. We can plot dominant eigenvalues of the system against to  $\beta$  to get some information about stability of the system according to changing of this parameter. Until  $\beta$  reaches a critical value, the norms are less than 1 and the system is stable. If  $\beta$  is increased beyond this critical value, the norms will be greater than 1 and stability of the system switches to unstable situation (see Figure 3). We can also determine this critical value of  $\beta$  by using the Schur-Cohn criterion that is given as follows.

**THEOREM 3.1 [7]** *A pair of complex conjugate roots of equation*

$$p(\lambda) = \lambda^3 + p_2\lambda^2 + p_1\lambda + p_0 \tag{17}$$

*lie on the unit circle and the other roots of equation (17) all lie inside the unit circle if and only if*

- (a)  $p(1) = 1 + p_2 + p_1 + p_0 > 0$  and  $p(-1) = 1 - p_2 + p_1 - p_0 > 0$ ,
- (c)  $D_2^+ = 1 + p_1 - p_0^2 - p_0p_2 > 0$ ,
- (d)  $D_2^- = 1 - p_1 - p_0^2 + p_0p_2 = 0$ .

Now, we return matrix  $\mathbf{J}_F(\bar{E})$  to determine bifurcation point of system (7). Computations give us that the exact characteristic polynomial (there is no assumption on the matrix) of matrix  $\mathbf{J}_F(\bar{E})$  is the form Equation (17) where

$$\begin{aligned} p_2 &= -1 - \exp(-K_1r_1\bar{M}) - \exp(-K_2r_2\bar{Z}), \\ p_1 &= \exp(-K_1r_1\bar{M}) + \exp(-K_2r_2\bar{Z}) + \exp(-K_1r_1\bar{M} - K_2r_2\bar{Z}) \\ &+ \bar{N} \frac{\beta^2 K_1 r_1 (1 - \exp(-K_2 r_2 \bar{Z})) - K_2 r_2 \alpha_1 \alpha_2 (1 - \exp(-K_1 r_1 \bar{M}))}{K_1 r_1 K_2 r_2}, \end{aligned}$$




Figure 4. Graph of Neimark-Sacker bifurcation of system (7) for  $\beta = 7.95907 \times 10^{-8}$ , where  $M(0) = 1 \times 10^6$ ,  $N(0) = 1.5 \times 10^5$ ,  $Z(0) = 9 \times 10^5$ . The other parameters are taken Example 2.3.

$$p_0 = -\exp(-K_1 r_1 \bar{M} - K_2 r_2 \bar{Z}) - \frac{\beta^2 \bar{N}}{K_2 r_2} \exp(-K_1 r_1 \bar{M})(1 - \exp(-K_2 r_2 \bar{Z})) + \frac{\alpha_1 \alpha_2}{K_1 r_1} \bar{N} \exp(-K_2 r_2 \bar{Z})(1 - \exp(-K_1 r_1 \bar{M})).$$

*Example 3.2* Solving equation c of Theorem 3.1, we have  $\bar{\beta} = 7.95907 \times 10^{-8}$ . Furthermore, we have also  $p(1) = 0.000132031 > 0$ ,  $p(-1) = 7.46124 > 0$  and  $D_2^+ = 0.500887 > 0$  for  $\bar{\beta}$ . Figure 4 shows that  $\bar{\beta}$  is the Neimark-Sacker bifurcation point of the system with eigenvalues  $\lambda_1 = 0.865769$ ,  $|\lambda_{2,3}| = |0.999508 \pm 0.0313588i| = 1$ , where blue, red and black graphs represent  $M(n)$ ,  $N(n)$  and  $Z(n)$  population densities, respectively.

#### 4. Result and discussion

In this paper, we have modified the tumour growth model proposed in [1] using system of differential equation with piecewise constant arguments that includes both discrete and continuous time for the populations. Some theoretical results for the boundedness, local and global stability of the system are obtained. The parameter values are selected from the study [1] which are obtained results of experiment on the dynamics of growth of highly malignant B Lymphoma Leukemic cells in the spleen of chimeric mice [1]. We observe that the parameter  $\alpha_1$  (decay rate of tumour cells) and parameter  $\beta$  (conversion rate from resting to hunting cells) play a key role to control the unlimited growth of tumour cells so as to control the oscillations of the tumour cells. Local stability analysis shows that if the parameter  $\alpha_1$  is less than a threshold value  $1.35777 \times 10^{-7}$  and parameter  $\beta$  is in interval  $4.2911 \times 10^{-9} < \beta < 1.0429 \times 10^{-7}$ , then tumour, hunting and resting cell coexist as a stable steady state (Figure 1). In addition, global stability analysis implies that the stability of the system with local stability conditions does not depend on initial conditions of the tumour, hunting and resting populations (Figure 2).

The Neimark-Sacker bifurcation is the discrete analogue of the Hopf bifurcation that occurs in continuous systems such as in [1]. In their study, a stable limit cycle constructed by Hopf

bifurcation is formed around the equilibrium point which depends on changing the parameter  $\tau$  and  $\beta$ . This result is also valid for system (7). As seen in Figure 4, stable periodic solutions occur at Neimark-Sacker bifurcation point (that is  $\bar{\beta} = 7.95907 \times 10^{-8}$ ) as a result of stable limit cycle. The existence of periodic solutions is relevant in tumour growth models. It means that the tumour population may oscillate around a equilibrium point even in the absence of any treatment. Such a phenomenon, which is known as Jeff 's Phenomenon, has been observed clinically [1]. When the value of parameter  $\beta$  is less than  $\bar{\beta}$  which falls in stable region (see Figure 3), the solution of system (7) has damped oscillation and the positive equilibrium point is local asymptotically stable (see Figure 5 for  $\beta = 2 \times 10^{-8}$ ). This implies that tumour, hunting and resting cell coexist as a stable steady state as a result of competition, namely tumour dormancy. If the value of parameter  $\beta$  is greater than  $\bar{\beta}$  which falls in unstable region (see Figure 3), the system has unstable oscillation and the positive equilibrium point is unstable (see Figure 6




Figure 5. Graph of iteration solution of the system for  $\beta = 2 \times 10^{-8}$ . The other parameters and initial conditions are the same as Figure 4.




Figure 6. Graph of iteration solution of the system for  $\beta = 1.5 \times 10^{-7}$ . The other parameters and initial conditions are the same as Figure 4.




Figure 7. Graph of the iteration solution of  $M(n)$ ,  $N(n)$  and  $Z(n)$ , where  $\alpha = 2.74379 \times 10^{-6}$ . The other parameters are the same as Figure 6.

for  $\beta = 1.5 \times 10^{-7}$ ). In this situation, tumour cells have growing oscillation with very high amplitude. We also investigate the dynamic behaviour of the system in the region ( $\beta > \bar{\beta}$ ) where the system exhibits unstable oscillation. In this region, decay rate of tumour cells  $\alpha_1$  plays an important role in controlling tumour cells growth. As the parameter  $\alpha_1$  is increased in this region, the population size of tumour cells can be constrained to null values namely tumour-free steady state where the tumour cells are eliminated by hunting cells. Therefore, it is possible to reach the tumour-free steady state by increasing the parameter  $\alpha_1$  (Figure 7).

When our theoretical and numerical results are compared with that of work [1], we obtain a good compatibility. Hence, differential equations with piecewise constant arguments may be used to approximate delay differential equations that contain discrete delays [3].

### Disclosure statement

No potential conflict of interest was reported by the authors.

### References

- [1] S. Banerjee and R.R. Sarkar, *Delay-induced model for tumor-immune interaction and control of malignant tumor growth*, Biosystems 91 (2008), pp. 268–288.
- [2] F. Bozkurt, *Modeling a tumor growth with piecewise constant arguments*, Discret. Dyn. Nat. Soc. (2013), Article ID 841764.
- [3] K.L. Cooke and I. Györi, *Numerical approximation of the solutions of delay differential equations on an infinite interval using piecewise constant arguments*, Comput. Math. Appl. 28 (1994), pp. 81–92.
- [4] K. Gopalsamy and P. Liu, *Persistence and global stability in a population model*, J. Math. Anal. Appl. 224 (1998), pp. 59–80.
- [5] F. Gurcan and F. Bozkurt, *Global stability in a population model with piecewise constant arguments*, J. Math. Anal. Appl. 360 (2009), pp. 334–342.
- [6] S. Kartal, *Mathematical modeling and analysis of tumor-immune system interaction by using Lotka-Volterra predator-prey like model with piecewise constant arguments*, PEN 2 (2014), pp. 7–12.
- [7] X. Li, C. Mou, W. Niu, and D. Wang, *Stability analysis for discrete biological models using algebraic methods*, Math. Comput. Sci. 5 (2011), pp. 247–262.
- [8] P. Liu and K. Gopalsamy, *Global stability and chaos in a population model with piecewise constant arguments*, Appl. Math. Comput. 101 (1999), pp. 63–88.

- [9] R.M. May, *Biological populations obeying difference equations: Stable points, stable cycles and chaos*, J. Theor. Biol. 51 (1975), pp. 511–524.
- [10] R.M. May and G.F. Oster, *Bifurcations and dynamic complexity in simple ecological models*, Am. Nat. 110 (1976), pp. 573–599.
- [11] Y. Muroya, *Persistence, contractivity and global stability in logistic equations with piecewise constant delays*, J. Math. Anal. Appl. 270 (2002), pp. 602–635.
- [12] Y. Muroya, *New contractivity condition in a population model with piecewise constant arguments*, J. Math. Anal. Appl. 346 (2008), pp. 65–81.
- [13] I. Ozturk and F. Bozkurt, *Stability analysis of a population model with piecewise constant arguments*, Nonlinear Anal. Real. 12 (2011), pp. 1532–1545.
- [14] I. Ozturk, F. Bozkurt, and F. Gurcan, *Stability analysis of a mathematical model in a microcosm with piecewise constant arguments*, Math. Biosci. 240 (2012), pp. 85–91.
- [15] J.W.-H. So and J.S. Yu, *Global stability in a logistic equation with piecewise constant arguments*, Hokkaido Math. J. 24 (1995), pp. 269–286.
- [16] K. Uesugi, Y. Muroya, and E. Ishiwata, *On the global attractivity for a logistic equation with piecewise constant arguments*, J. Math. Anal. Appl. 294 (2004), pp. 560–580.